

## REAL TIME PCR DIAGNOSIS OF LEISHMANIASIS IN CONSTANTINE (ALGERIA)

Reçu le 13/03/2011 – Accepté le 15/06/2012

MIHOUBI I. <sup>a</sup>, FRAHTIA K. <sup>a</sup>, GASSEM N. <sup>b</sup>, PICOT S. <sup>c</sup>, DEMONBRISON F. <sup>c</sup>

<sup>a</sup>Laboratoire de Biosystématique et Ecologie des Arthropodes, Université Mentouri, Constantine, Algérie

<sup>b</sup>Laboratoire de Parasitologie – Mycologie Médicale, CHU Constantine, Algérie

<sup>c</sup>Service de Paludisme, Parasites du sang et Mycologie Médicale, Hôpital de la Croix-Rousse, Hospices civils de Lyon, Lyon, France

### Résumé

Compte tenu de la réémergence de la leishmaniose dans le monde, de la variabilité géographique de l'épidémiologie et de l'augmentation du nombre de voyageurs, une étude pilote sur le diagnostic des leishmanioses cutanée et viscérale a été réalisée à Constantine, l'un des foyers de l'Est algérien. 143 prélèvements cutanés et 108 échantillons de moelle osseuse ont été récoltés sur buvards et testés par PCR en temps réel et les résultats comparés à ceux de l'examen direct. Le diagnostic de la LC par PCR a été positif dans 81% des cas pour la PCR contre 48% pour la microscopie. Quant à la LV, le taux de positivité est de 29,6% pour la PCR en temps réel contre 24% pour la microscopie. La PCR en temps réel a montré une différence quantitative significative entre les patients pour lesquels le diagnostic microscopique a été positif et ceux dont l'examen direct a été négatif. Les résultats présentés dans cette étude ont montré l'efficacité et la sensibilité de la PCR dans le diagnostic des leishmanioses cutanée et viscérale à partir des buvards. Avec cette technique, il a été possible de réaliser sur place un prélèvement à chaque malade et d'obtenir des résultats dans un délai court. Une collaboration Nord-Sud basée sur l'utilisation de moyens simples de transmission des échantillons pour le diagnostic moléculaire a permis de créer un partenariat efficace en terme de diagnostic quotidien et favoriser ainsi l'échange de chercheurs afin de préparer le transfert de technologie.

**Mots clés :** *Leishmaniose cutanée – Leishmaniose viscérale - Diagnostic - PCR en temps réel - Algérie.*

### Abstract

Taking into account the re-emergence of leishmaniasis in the world, the geographic variability of its epidemiology and the growing numbers of travellers, a pilot study on the diagnosis of cutaneous leishmaniasis and visceral leishmaniasis was undertaken in Constantine, one of the outbreak regions in eastern Algeria. A total of 143 cutaneous specimens and 108 bone marrow samples were collected on blotters and tested by real-time PCR. Results were compared with those of direct examination. Diagnosis, in CL, was positive for leishmaniasis in 81% of cases using PCR versus 48% of cases using microscopy. For VL, results were 26.9% for real time PCR versus 24% for microscopy exam. Real-time PCR showed a significant quantitative difference between patients for whom microscopic diagnosis was positive and those for whom direct examination was negative. The results presented in this study demonstrated the effectiveness and sensitivity of PCR in the diagnosis of cutaneous and visceral leishmaniasis from blotter specimens. This technique enabled in-field collection of specimens from each patient and provided prompt results. North- South cooperation based on the use of simple means for transmission of specimens for molecular diagnosis allowed creation of an effective partnership for daily diagnosis and promoted exchange between investigators in preparation for technology transfer

**Keywords:** *Cutaneous leishmaniasis – Visceral leishmaniasis – Diagnosis – Real time PCR- Algeria.*

### ملخص

نتيجة لعودة ظهور داء اللشمانيات في العالم ، للتغيرات الجغرافية من حيث الإصابات وزيادة عدد المسافرين ، تمت دراسة لتشخيص هذا المرض الجلدي و الحشوي بقسنطينة التي تعتبر إحدى البؤر بالشرق الجزائري. تمت تحليل 338 عينة جلدية و 112 عينة من مخ العظام بواسطة المجهر و PCR في الوقت الحالي. النتائج المحصل عليها بينت بان عدد المرض ذو نتيجة ايجابية يقدر بـ 81% بالنسبة لـ PCR في الوقت الحقيقي و 48% بالنسبة لتشخيص المجهرى لليشمانيا الجلدية. أما داء اللشمانيات الحشوي، أظهرت PCR في الوقت الحقيقي النتائج المحصل عليها : 29,6% PCR في الوقت الحقيقي، مقابل 24 % للفحص المجهرى. وأظهرت PCR في الوقت الحقيقي فرق كبير بين المرضى الذين الكمية التشخيص المجهرى كان ايجابيا وأولئك الذين الفحص المباشر كان سلبيا. وأظهرت النتائج الواردة في هذه الدراسة فعالية بكر في تشخيص اللشمانيا الجلدية والحشوية من السجلات اليومية. مع هذا الأسلوب كان من الممكن جمع العينات لكل مريض والنتائج في وقت قصير وإنشاء تعاون بين الشمال والجنوب على أساس استخدام وسائل بسيطة لنقل العينات للتشخيص الجزيئي شراكة فعالة من حيث التشخيص وبالتالي تشجيع تبادل يوميا من الباحثين لإعداد ونقل التكنولوجيا.

**الكلمات المفتاحية :** *الليشمانيات الجلدي -- الليشمانيات الحشوي - التشخيص-PCR في الوقت الحقيقي --الجزائر.*



Leishmaniasis is a parasitic infectious disease comprised from a variety of syndromes, which are different between them in pathogenesis and clinical picture. According to the World Health Organization (WHO), leishmaniasis is considered as one of the most important parasitic diseases [1]. It is endemic in 88 countries in five continents with 12 million people afflicted world wide from it and another 350 million living at risk of infection. About 1-1.5 million new cases of cutaneous leishmaniasis (CL) and 500,000 of visceral leishmaniasis (VL), rise per year [2].

Algeria people, affected by two kinds of leishmaniasis; cutaneous leishmaniasis (CL) and visceral leishmaniasis (VL).

Cutaneous leishmaniasis which is a serious public health problem in Algeria. It has two clinical forms caused by *Leishmania major* and *L. infantum* [3] [4]. *Leishmania infantum* zymodeme MON-24 is the main strain responsible for the sporadic CL distributed throughout the northern part of the country [5]. The vector is *Phlebotomus perfiliewi* [6] and dogs constitute the most likely reservoir [7]. *Leishmania major* zymodeme MON-25 is present in the steppe areas of central and southern parts of the country. *Psammomys obesus* and *Meriones shawi* are its proven reservoir hosts [8] [9] and *P.papatasi* is the main vector [10].

Visceral leishmaniasis due to *L. infantum* is the most sever form. It is fatale if untreated. It occurs mainly in the north where it reaches almost 540 per year. The incidence of visceral leishmaniasis is 0.31 cases per 100,000 inhabitants [11]. The outbreak of the Great Kabylia alone accounts for nearly 50% of cases [12] where the disease is responsible for 6% of deaths [13].

Diagnosis of this disease, in Algeria, is still based on direct examination and *in vitro* culture of bone marrow aspirates or detection of antibodies by immunofluorescence [14].

In recent years, real-time polymerase chain reaction (PCR) has been developed and used successfully for the diagnosis of leishmaniasis [15][16]. This procedure is non-invasive, highly sensitive and specific, and useful in monitoring responses to therapy by quantification of parasitic load [17]. The aim of the present study was to propose a specific real-time PCR approach for the diagnosis of the two forms of leishmaniasis imprinted onto filter paper.

## MATERIAL AND METHODS

### Parasites samples

#### *Leishmania* reference strains

*L. infantum* MHOM/TN/80/IPT1, *L. major* MHOM/SU/73/5-ASKH were kindly provided by M. Gramiccia, Istituto Superiore di Sanita in Roma, Italy. The strains were cultivated at 25 °C in RPMI medium

(Invitrogen, Cergy Pontoise, France) supplemented with 15% foetal calf serum and Penicillin. Parasites were harvested at a density of 10<sup>4</sup> parasites/ml. They were washed twice with PBS buffer, pelleted by centrifugation (5900×g for 3 min) and stored at -20 °C.

## Human clinical investigations

### Cutaneous lesions

A total of 143 skin lesions were collected from 143 patients consulting at the parasitological laboratory in the east of Algeria (Constantine Hospital) from the period of September 2004 to May 2006.

The CL are the typical lesions skin not affected with fungi and bacteriological infection and the absence of specific treatment during three months before the consultation. All individuals enrolled in the study provided informed consent.

**Bone Marrow samples:** A total of 108 bone marrow samples were collected on filter paper 3MM Watmann From children with visceral leishmaniasis in the east of Algeria (Constantine hospital) from the period of 2004 until 2006, on the basis clinical symptoms (anarchic fever, splenomegaly, hepatomegaly, mucocutaneous and pallor) and biological arguments (normocytic anemia, pancytopenia, thrombocytopenia...).

Cutaneous lesions and bone marrow samples have been spotted on 3mm Whatman™ filter paper, air-dried and stored in individual plastic zip-lock bag at room temperature.

All individuals enrolled in the study provided informed consent.

## Nucleic acid isolation

### *Leishmania* reference strains

DNA was extracted using QiAamp DNA mini kit (Qiagen, Courtaboeuf, France) according to the manufacturer's instructions with minor modifications. The incubation time with proteinase K was increased to 1 h at 56 °C and DNA was eluted from the column with 50 µl of PCR-grade H<sub>2</sub>O.

### Filter paper samples

The DNA was isolated from the dried cutaneous spots using Instagene Matrix® resin (Bio-Rad, Marnes la Coquette, France). A 1-cm paper circle was cut avoiding the inclusion of non-impregnated filter paper. Paper disks were cut in small sectors in a sterile Eppendorf tube and soaked in Saponin 5% during 60 min. The red residue was collected by centrifugation and papers were incubated with 200µl Instagene Matrix® resin at 56 °C for 30 min with vortexing after 15 min and at the end of incubation. Tubes were transferred in a boiling bath for 8 min before



centrifugation and supernatants were decanted carefully to avoid resin carry-over. The extracted DNA was diluted in sterile water to 1/5 and kept at  $-20^{\circ}\text{C}$  before PCR.

### Polymerase chain reaction amplification

Each sample has been subjected to real-time PCR assay for the detection of *Leishmania* DNAs as previously described [15]. The samples presenting a crossing point (point which marked the cycle at which the fluorescence of the sample is significantly different from the baseline signal) lower or equal to 25 have been selected for the identification of old world *Leishmania* species. The primers JW11 (5'-CCTATTTTACACCAACCCCAAGT-3') and JW12 (5'-GGGTAGGGCGTTCTGCGAAA-3'), designed on the conserved region of *Leishmania* kDNA minicircle, were used to allow the amplification of the genus *Leishmania* [18]. Real-time PCR technology was performed with fluorescent SYBR Green I. The extracted DNA (5  $\mu\text{l}$ ) was added to 15  $\mu\text{l}$  of reaction mixture containing 4mM of  $\text{MgCl}_2$ , 0.5  $\mu\text{M}$  of each primer (JW11 and JW12), 2  $\mu\text{l}$  of LightCycler FastStart DNA Master SybrGreen I buffer (Roche Applied Science, Meylan, France). Conditions for cycling were  $95^{\circ}\text{C}$  for 4 min, followed by 35 cycles of  $95^{\circ}\text{C}$  for 10 s,  $62^{\circ}\text{C}$  for 10 s, and  $72^{\circ}\text{C}$  for 10 s with monitoring of fluorescence at the end of elongation step. PCR product identity was confirmed with melting curve analysis. Melting program consists of one cycle of  $95^{\circ}\text{C}$  for 0 s,  $67^{\circ}\text{C}$  for 20 s and heating at  $98^{\circ}\text{C}$ . The transition rates were  $20^{\circ}\text{C}/\text{s}$  except for the extension step and the final step, which have a temperature transition rate at  $1^{\circ}\text{C}/\text{s}$  and  $0.1^{\circ}\text{C}$ , respectively.

An uninfected sample was systematically included to control the amplification specificity. DNA isolated from the *Leishmania* reference strains has been used as a positive control in each run. The sensitivity of the PCR assay was assessed on a serial ten fold dilutions ranging from 1000 to 1 parasite/ml of DNA purified from the four *Leishmania* references strains. The detection limit was 10 parasite /ml corresponding to 0.05 parasites per reaction (data not shown).

## RESULTS

### Human clinical investigations

#### Cutaneous lesions

During this study, 143 patients were included. The sex ratio is 1.55 (87 men/56 women). The mean ages of men patients were  $28.3 \pm 16.5$  years and for women were  $35 \pm 8.5$ . Children (under 16 years) account for 17.5% of all patients. Clinically, the lesions were most frequently on exposed parts: faces (61 patients: 45%) and members (42 patients: 31%). The lesions were multiple and sometimes showed different aspects. They ranged between 1 and 4 cm in diameter.

#### Bone marrow samples

In this present study 108 patients were included. The average age of children was from  $7.20 \pm 3.53$  years (males:  $4.86 \pm 4.43$  years, girls:  $5.01 \pm 4.28$  years) and sex ratio of 1.3.

For both types of leishmaniasis, patients were mostly from Constantine and its surroundings (Ibn Ziad, Hamma Bouziane, Ain Abid, Chelghoum Laid).

### Molecular diagnosis

#### Cutaneous samples

The real-time PCR was positive for 117/143 samples (81.81%). The increased sensitivity of laboratory diagnosis obtained by the use of PCR compared to the direct examination was therefore 46/143 patients (32,16%). All patients with a negative result by PCR had a negative microscopic examination.

The real-time PCR by Light Cycler, showed that there was a significant quantitative difference between patients whose diagnosis was microscopically positive and those who were negative. Indeed, patients for whom microscopic examination was positive showed a detection limit of  $19.56 \pm 0.108$  average PCR cycles with an average of  $28.47 \pm 0.059$  ( $p < 0.01$ ) in patients whose direct examination was negative.

#### Bone marrow samples

The real time PCR was positive for 32/108 samples, representing 29.6% of all samples analyzed. Cycles of positivity were observed in average of 25.77. In all positive cases, the denaturation curves showed the expected characteristics of specificity.

The results of molecular diagnosis were compared with those obtained by microscopy. It shows that increasing the sensitivity of laboratory diagnosis, obtained by using PCR, compared to the direct examination was 6/108 patients (5.5%). In fact, 6 patients were negative on microscopic examination and positive by real-time PCR. The positive rate of direct examination was about 24%.

## DISCUSSION

The yearly incidence of leishmaniasis, estimated at 1.5 millions for CL and 0,5 million for LV, is increasing related to deforestation, urbanization or environmental changes [18].

Given its reemergence in Algeria, it was important to establish a pilot study on the diagnosis

of both forms of the disease [19]. Recent data show that the incidence is increasing, and geographic range extension [20] [14].

Visceral leishmaniasis takes, with cutaneous leishmaniasis, the first place in parasitic diseases and accounts for 35% of notifiable diseases [21]. This requires a rapid and sensitive diagnosis.



The current diagnosis is usually based on direct examination and culture which are difficult because of low parasite loads of samples, and difficult to achieve under experimental conditions that do not always fulfil all necessary conditions. Obviously, as with most parasitic diseases, the contribution of molecular diagnostics can be considerable [22][23] [24] [25] [26][27]. PCR is a diagnosis method confirmed cutaneous leishmaniasis [15] or mucocutaneous leishmaniasis [28].

The real-time PCR, as new technology has revolutionized molecular diagnostics balancing reliability and timeliness [29]. Indeed, early diagnosis brings PCR can identify leishmaniasis before the clinical picture is complete (suspect). The skin and bone marrow samples are very suitable for detection of *Leishmania* DNA by PCR [30].

Several authors report its interest in the diagnosis of leishmaniasis [31].

It was used in the diagnosis of VL of the child in several Mediterranean countries [16]. It has also shown its benefits in the detection of asymptomatic carriers of the parasite in patients infected with HIV for whom, VL coinfection is frequently reported in countries where both infections (cutaneous and visceral) are endemic [32]. It has proved useful in monitoring therapy by quantification of parasite load [16].

It can be applied to different types of samples from reservoirs and vectors, hence its interest in epidemiological studies [33]. Quantitatively, the RT-PCR seems to be the most appropriate in evaluating the success of therapy and new types of treatments [34].

Moreover, the collection of bone marrow samples on filter paper is a good alternative for preserving samples for molecular analysis relocated to a reference laboratory. Its use in the diagnosis of cutaneous leishmaniasis and the identification of different species of *Leishmania* has been applied with success [35] [36]. In our study, we included all patients with clinical and biological signs suggestive of visceral and cutaneous leishmaniasis.

The real-time PCR increased the number of certain documentation of diagnosis from 71 to 117 cases for CL and 26 to 32 cases for VL, in a suspected population with respectively, 143 and 108 patients. As part of this retrospective study, it was not possible to follow the clinical evolution of patients and we have no information to confirm absolutely that patients identified have visceral leishmaniasis and that treatment made a specific improvement or even a cure. However, it was not the aim of the study show that the real-time PCR to better care for patients, but rather to demonstrate the concept that this technique, applied to samples preserved on blotting paper and easily transportable, can give better results in terms of detection rate of visceral leishmaniasis.

This concept is important since it may allow institutions to mobilize the country to promote the establishment of reference laboratories equipped with real-time PCR technology. The availability of this powerful tool will lead the development of clinical and biological studies more complete systems that provide definitive answers to the conditions for improving the diagnosis of this severe disease.

## REFERENCES

- 1- WHO (1990) Control of the Leishmaniases. Technical Report Series 793. WHO, Geneva, Switzerland, pp. 1–158.
- 2- Desjeux P., Alvar J., Gradoni L., Gramiccia M., Medrano F.J., Deniau M. et al. Epidemiological analysis of 692 retrospective cases of *Leishmania*/HIV co-infections. *WHO/LEISH*, 1996 ; 39: 1-11
- 3- Belazzoug S. Les leishmanioses en Algérie. A propos de l'identification de 32 souches d'origine humaine et animale. *Leishmania*. Taxonomie et phylogénèse. Applications éco-épidémiologiques (Coll. Int. CNRS/INSERM, 1984). Montpellier: IMEEE; 1986a. p. 397-400
- 4- Harrat Z, Pratlong F, Belazzoug S, Dereure J, Deniau M, Rioux JA, et al. *Leishmania infantum* and *Leishmania major* in Algeria. *Trans R Soc Trop Med Hyg* 1996; 90:625-629.
- 5- Belazzoug S, Ammar-Khodja A, Belkaid M, Tabet-Derraz O. Cutaneous leishmaniasis in northern Algeria [in French]. *Bull Soc Pathol Exot Filiales* 1985;78:615- 622.
- 6- Izri MA, Belazzoug S. *Phlebotomus (Larroussius) perfiliewi* naturally infected with dermatotropic *Leishmania infantum* at Tenes (Algeria). *Trans R Soc Trop Med Hyg* 1993;87:399.
- 7- Benikhlef R, Harrat Z, Toudjine M, Djerbouh A, Bendali-Braham S, Belkaid M. Présence de *Leishmania infantum* MON-24 chez le chien. *Méd Trop* 2004;64:42-46.
- 8- Belazzoug S. Isolation of *Leishmania major* Yakimoff & Schokhor, 1914 from *Psammomys obesus* Gretzschmar, 1828 (Rodentia: Gerbillidae), in Algeria. *Trans R Soc Trop Med Hyg* 1983;77:876.
- 9- Belazzoug S. Découverte d'un *Meriones shawi* (Rongeur, Gerbillidé) naturellement infesté par *Leishmania* dans le nouveau foyer de leishmaniose cutanée de K'sar Chellala (Algérie). *Bull Soc Pathol Exot* 1986;79:630-633.
- 10- Izri MA, Belazzoug S, Pratlong F, Rioux JA. Isolation of *Leishmania major* in *Phlebotomus papatasi* in Biskra (Algeria). The end of an ecoepidemiological saga [in French]. *Ann Parasitol Hum Comp* 1992; 67:31-32.





- 11- Institut National de Santé Publique. Ministère de la Santé de la Population et de la Réforme Hospitalière. Direction de la Prévention. Relevé annuel des maladies à déclaration obligatoire Année 2005. Alger: Institut National de Santé Publique; 2005. <http://www.sante.dz/insp/rem.html> [accessed 16 April 2009].
- 12- Harrat Z., Addadi, K., Tabet Deraz O. - La leishmaniose viscérale en Algérie : recensement des cas de leishmaniose viscérale. *Bull Soc Path Exo* 1992; 85: 296-301.
- 13- Djerbouh A., Toudjine M., Djoudi M., Benikhlef R., Harrat Z. - La leishmaniose canine en Algérie : essai de traitement par l'allopurinol. *Ann Méd Vét* 2005 ; 149 : 132-134.
- 14- Izri A. ; Belazzoug S. – Diagnostic de laboratoire des leishmanioses rencontrées en Algérie. *Revue francophone des laboratoires* 2007 ; Supplément au numéro 396,
- 15- Mihoubi I., De Monbrison F., Romeuf N., Moulahem T., Picot S. - Diagnostic délocalisé par PCR en temps réel de la leishmaniose cutanée dans le foyer de Constantine (Algérie). *Med Trop* 2006; 66: 39-44
- 16- Aoun K., Chouih E., Amri F., Ben Alaya N., Raies A., Mary C., Bourabrine A. Short Report: Contribution of Quantitative Real-Time Polymerase Chain Reaction to Follow-Up of Visceral Leishmaniasis Patients Treated with Meglumine Antimoniate. *Am. J. Trop. Med. Hyg.*, 2009; 81(6): 1004-1006
- 17- Mary C, Farault F, Droguoul MP, Xeridat B, Scleinitz N, Cuisenier, Dumon H., Reference values for *Leishmania infantum* parasitemia in different clinical presentations: quantitative polymerase chain reaction for therapeutic monitoring and patient follow-up. *Am J Trop Med Hyg*, 2006; 75: 858–863.
- 18- Nicolas, L., Milon, G., Prina, E., 2002. Rapid differentiation of old world *Leishmania* species by LightCycler polymerase chain reaction and melting curve analysis. *J. Microbiol. Methods*. 51,295–299.
- 19- Desjeux P. The increase in risk factors for leishmaniasis worldwide. *Trans R Soc Trop Med Hyg* 2001;95:239-243.
- 20- Mott KE, Nuttall I, Desjeux P, Cattand P. New geographical approaches to control of some parasitic zoonoses. *Bull World Health Organ*. 1995;73(2):247-57.
- 21- Barchiche N.A., Madiou M. - Recrudescence des leishmanioses cutanées : à propos de 213 cas dans la wilaya de Tizi-Ouzou. *Pathol. Biol*. 2009 ; 57 : 65–70
- 22- Reithinger R., Dujardin J.C.- Molecular diagnosis of leishmaniasis: current status and future applications. *J. Clin. Microbiol*. 2007; 45:21–25.
- 23- Alam M.Z., Abul Khair. Shamsuzzaman M., Kuhls K. , Schönian G. - PCR diagnosis of visceral leishmaniasis in an endemic region. Mymensingh district, Bangladesh. *Trop. Med, International Health* 2009; 14 (5) : 499–503
- 24- Deborggraeve S., Boelaert M., Rija S., De Doncker S., Dujardin J.C., Herdewijn P., Bütscher P. - Diagnostic accuracy of a new *Leishmania* PCR for clinical visceral leishmaniasis in Nepal and its role in diagnosis of disease. *Trop. Med, International Health* 2008; 13 (11): 1378–1383
- 25- Pandey K., Pandey B.D., Mallik A.K., Kaneko O., Uemura H., Kanbara H., Yanagi T., Hirayama K. - Diagnosis of visceral leishmaniasis by polymerase chain reaction of DNA extracted from Giemsa's solution-stained slides. *Parasitol Res* 2010 ; 107:727–730
- 26- Souza Leite R., de Almeida Ferreira S., Trindade Ituassub S., de Melob M.N., Silva Ribeiro de Andrade A. - PCR diagnosis of visceral leishmaniasis in asymptomatic dogs using conjunctival swab samples. *Vet Parasitol* 2010;170 : 201–206
- 27- Fraga Leite T., Brustoloni Y., M., Lima R.B., Cavalheiros Dorva M.E., Oshiro E.T., Oliveira J., Lyrio De Oliveira A., Pirmez C. - Polymerase chain reaction of peripheral blood as a tool for the diagnosis of visceral leishmaniasis in children. *Mem Inst Oswaldo Cruz*, Rio de Janeiro 2010; 105(3): 310-313.
- 28- Nasereddin A., Benssoussan- Hermano E., Schönian G., Baneth G., Jaffe C.L. - Molecular Diagnosis of Old World Cutaneous Leishmaniasis and Species Identification by Use of a Reverse Line Blot Hybridization Assay. *J.Clin. Microbiol* 2008; 2848–2855.
- 29- Bastien P, Procop G.W., Reischl U. - Quantitative real-time PCR is not more sensitive than "conventional" PCR. *J Clin Microbiol*. 2008, 46(6):1897-1900.
- 30- Quaresma P.F., Fonseca Murta S.M., DE Castro Ferreira E., Mariano DA Rocha-Lima A.V., Prates Xavier A.A., Ferreira Gontijo C.M. Molecular diagnosis of canine visceral leishmaniasis: Identification of *Leishmania* species by PCR-RFLP and quantification of parasite DNA by real-time PCR. *Acta Tropica* 2009; 111 ; 289-294.
- 31- Schönian G, Kuhls K, Mauricio IL. Molecular approaches for a better understanding of the epidemiology and population genetics of *Leishmania*. *Parasitology*, 2010; 16:1-21
- 32- Neffati A., Kallel K., Anene S., Kaouech E., Belhadj S., Ennigrou S., Chaker E. Choix des amorces : élément déterminant dans le diagnostic moléculaire de la leishmaniose cutanée. *Pathol Biol*, 2009, [Epub ahead of print]



33- Silva E.S, Gontijo CMF, Melo MN - Contribution of molecular techniques to the epidemiology of neotropical *Leishmania* species. *Trends in Parasitol* 2005; 21 : 550-552.

34- Frank D.T., Nasereddin A., Schnur L.F., Schonian G., Toz S.O., Jaffe C. L., Baneth G. Detection and Identification of Old World *Leishmania* by High Resolution Melt Analysis. *Neglected Tropical diseases* 2010; 4: Issue 1, e581.

35- De Monbrison F., Angei F., Staal A., Kaiser K., Picot S. - Simultaneous identification of the four human plasmodium species and quantification of plasmodium DNA load in human blood by real-time PCR. *Trans R Soc Trop Med Hyg* 2003; 97: 1- 4.

36- Mihoubi I., Picot S., Hafirassou N.; Monbrison de F. - Cutaneous leishmaniasis caused by *Leishmania tropica* in Algeria. *Trans R Soc Trop Med Hyg* 2008; 102, 1157-1159

